

Research Review

EDUCATIONAL SERIES™

Antimicrobial Mouthrinse in Dental Practice and Home Oral Hygiene Smoking Cessation and Oral Health

About the Experts



Vivian Chan Virata
DDM, MS, FADI, FIGD, FPFA, FPAGD

Dr. Vivian Virata works in private practice in Manila where she is Head of Periodontics Department at the Asian Hospital and Medical Center. She obtained her undergraduate degree in Dental Medicine from the University of the Philippines, and her postgraduate Master's of Science degree in Periodontology from Columbia University in New York. Since 2001 she has been a Professorial lecturer in Periodontics at the College of Dentistry at the University of the Philippines. She is Vice President of the Philippine Society of Periodontology and Past President of the Philippine Academy of General Dentistry. Dr Virata has been invited to speak at many conferences.



Professor Edward C. M. Lo
BDS, MDS, PhD, FHKAM, FCDSHK

Dr. Edward Lo is the Chair Professor of Dental Public Health in the Faculty of Dentistry of the University of Hong Kong and is a guest professor in a number of major dental schools in mainland China. He has obtained his full academic and specialist training in Hong Kong, including BDS and PhD degrees, and Fellowship of the Hong Kong Academy of Medicine. Prof. Lo is the current Chairman of the Specialty Board of Community Dentistry of the College of Dental Surgeons of Hong Kong. He is a past President of the International Association of Dental Research (IADR) Southeast Asian Division and that of the Asia-Pacific Region. His expertise is in oral epidemiology, oral health care delivery and preventive dentistry. He has published over 200 papers in international scientific journals.

ABOUT RESEARCH REVIEW

Research Review is an independent medical publishing organisation producing electronic publications in a wide variety of specialist areas. Research Review publications are intended for health care professionals. Our publications range from regular updates of medical literature to synopses of speaker events and conferences, as well as individually commissioned pieces focused on specific disease states or medications.

Content is created independently of sponsor companies with assistance from leading local specialists.

Publications are free to receive for health care professionals, keeping them up to date with their chosen clinical area.

Subscribe free at www.researchreview.com/Philippines

Please send us any feedback or comments to admin@researchreview.com



This article is intended as an educational resource for dental healthcare professionals. It provides an overview of the following topics: 1) infection risk and control within the dental practice environment; 2) role of antimicrobial mouthrinses in infection control and patient oral hygiene; and 3) encouraging smoking cessation in smokers and facilitating access to smoking cessation programmes.

Infection risk in the clinic

Dental patients and dental healthcare professionals can be exposed to pathogenic micro-organisms in the dental setting, including bacteria and viruses, which can be transmitted via:¹⁻³

- Direct contact with blood, oral fluids, or other patient biomaterials.
- Indirect contact with contaminated objects or surfaces (e.g., charts, instruments, equipment).
- Contact of conjunctival, nasal, or oral mucosa with droplets generated from an infected person and propelled a short distance (e.g. by coughing, sneezing, or talking).
- Inhalation of airborne droplets that remain suspended in the air for long periods.

Aerosols, sprays, and splatter generated during routine dental procedures, especially during ultrasonic and air turbine procedures, can contain blood and saliva.^{3,4} The terms aerosols, sprays, and splatter are often used interchangeably to describe droplet particles; however, they differ in terms of their size. Mist-like aerosols are typically invisible and can remain airborne for prolonged periods of time. Splatter and spray consist of larger droplet particles, which can travel further than aerosols to land on the skin and other surfaces.³

Dental and oral health practitioners strive to manage these generated aerosols, sprays, and splatters by using personal protective equipment, barriers, and infection control protocols. However, practitioners may not fully appreciate that the spread of potential pathogenic micro-organisms is greater than previously considered and may encompass the majority of the dental operatory area (**Figure 1**).⁵ The behaviour of these droplet particulates and their associated health risks are complex,⁴ but aerosols, sprays, and splatters contaminated with pathogenic micro-organisms represent a potential route for disease transmission.^{3,6,7} Whether or not the spread of micro-organisms results in clinical infection depends in part on the virulence (infectivity) and dose (load) of a particular micro-organism and on the susceptibility of the host.^{3,7}




Figure 1. Mean number (with standard deviations of total counts) of colony-forming units (CFU) of different types of bacteria at various distances from treatment units after 1.5- and 3-hour collection times.⁵ Significant contamination was detected at all distances sampled when high-speed instruments were used.

Infection control in the clinic

A survey of dentists in seven Asian countries (including Thailand and the Philippines) conducted in the early 2000s suggested that knowledge, attitudes, and the practice of infection control and safety varied across the region and indicated a need for improved dental safety education and practice.⁸ More recently, a 2014 review of international studies of compliance with infection prevention and control in oral healthcare practices also suggests the need for greater infection prevention and control knowledge and education in developing countries.⁹

The purpose of infection control in dental practice is to prevent the transmission of pathogenic micro-organisms between patients and between dental staff and patients. The following procedures can minimize the spread of micro-organisms in the dental clinic setting:^{1,7}

- Use of personal protective equipment, including gloves, masks, and protective eyewear.
- Use of a high-volume evacuator, which exhausts externally during aerosol-creating procedures, such as ultrasonic and air turbine procedures.
- Use of a rubber dam to reduce the risk of contamination by infective aerosols (use whenever possible to isolate an area of the patient's mouth during treatment).
- Use of an antimicrobial mouthrinse by the patient prior to any intra-oral procedure, especially high-speed instrumentation – to reduce the micro-organism load in aerosols and splatter.

In addition to the routine use of personal protective equipment,^{1,7} the use of pre-procedural mouthrinses, high-volume evacuation, and rubber dam are the most effective methods of minimising the risk of exposure.⁴ Precautions such as these should be employed as standard practice because patients with bloodborne infections can be asymptomatic or unaware they are infected.¹

Precautions for infectious patients

Given that most of the procedures used in dentistry generate aerosols, patients with active infectious diseases (e.g. influenza) who require urgent dental treatment pose a considerable infection risk to dental staff and other patients.⁷ In such cases, the specific transmission-based precautions that must be followed include: scheduling these patients at the end of the day; use of pre-procedural antimicrobial mouthrinses and rubber dam; minimizing the use of aerosol-generating techniques; and applying two cycles of cleaning for environmental surfaces.

Pre-procedural mouth rinsing

The use of antimicrobial mouthrinses by patients prior to a dental procedure is intended to reduce the number of micro-organisms released from a patient in the form of aerosols or splatter that might contaminate a dental surgery and its equipment surfaces.¹ Pre-procedural rinsing may also reduce the number of micro-organisms accessing the patient's bloodstream during an invasive dental procedure.

There is no conclusive published evidence that pre-procedural mouth rinsing prevents clinical infection in dental staff or patients.¹ Nevertheless, many clinical studies have demonstrated that pre-procedural rinsing with essential oils-, chlorhexidine gluconate-, or cetylpyridinium chloride-based mouthrinses, either alone or together with use of a high-volume evacuator, is effective in reducing the microbial load of the aerosols produced during ultrasonic scaling.¹⁰⁻¹⁶

In one double-blind, randomised, cross-over study of patients undergoing ultrasonic scaling, pre-rinsing for 30 seconds with an essential oils-based mouthrinse resulted in a significant ($p < 0.001$) reduction in the number of colony-forming units in recoverable aerosol samples (Figure 2).¹⁴ Reduced bacterial load implies reduced risk of infection.




Figure 2. Percent reduction in colony-forming units (CFUs) contained in aerosols generated during 10-minute ultrasonic scaling performed after first rinsing with an essential oils-containing mouthrinse in a randomised, double-blind, cross-over study.¹⁴

PRACTICE TIPS 1 – INFECTION CONTROL

1. Dental healthcare professionals should be aware of the risk of disease transmission during dental procedures, especially when treating patients with active infectious diseases.
2. Aerosols and splatter generated during high-speed dental procedures are the primary means of potential disease transmission.
3. The following techniques should be employed to minimize the risk of exposure to aerosols and splatter:
 - I. Use of standard personal protective equipment.
 - II. Use of high-volume extraction.
 - III. Use of a rubber dam.
 - IV. Use of patient pre-procedural antimicrobial mouthrinse.
4. Seeing patients with active infectious disease at the end of the day.

Antimicrobial mouthrinses

The oral cavity harbours a vast variety of species of bacteria, viruses, and fungi, but it is bacteria that are the primary cause of periodontal disease.^{17,18} More than 300 species of bacteria associated with periodontal disease have been isolated from the oral cavity. Periodontal disease results from the establishment of dental plaque biofilm, which involves bacteria attaching to one of several oral surfaces, including the tooth and epithelium, as well as with other bacteria already attached to these surfaces.

Mechanical plaque biofilm removal through tooth-brushing and flossing is the gold standard for the prevention of periodontal disease and dental caries.¹⁹ However, most people fall short of optimal oral hygiene.^{20,21} Hence, the use of an antimicrobial mouthrinse is an important adjunct to professional care and tooth-brushing and flossing in the home. The most commonly used antimicrobial agents in clinical mouthrinses are: chlorhexidine gluconate, essential oils, and cetylpyridinium chloride.

Active ingredients and mechanisms of action

To varying degrees, chlorhexidine gluconate, essential oils, and cetylpyridinium chloride all disrupt the integrity of the bacterial cell membrane, leading to lysis and death (Table 1).^{21,22} The major advantage of chlorhexidine gluconate is its ability to bind (12-hour substantivity) to soft and hard oral tissues, enabling it to act over a long period after use and to inhibit adsorption of bacteria onto oral surfaces. Cetylpyridinium chloride binds to teeth and plaque to a lesser degree than chlorhexidine gluconate and is generally less efficacious than chlorhexidine gluconate. Chlorhexidine gluconate and essential oils penetrate plaque biofilm and produce changes in microbial cell surface morphology that alter co-aggregation, decolonisation, and, thus, survival.

| Active Ingredient | Description | Mechanism of action |
|--------------------------|---|--|
| Essential oils | Fixed combination of: Eucalyptol (0.092%) Menthol (0.042%) Methyl salicylate (0.060%) Thymol (0.064%) | Ruptures bacterial cell wall, leading to leakage of contents and cell death Penetrates the plaque biofilm to exert antimicrobial effects |
| Cetylpyridinium chloride | Quaternary ammonium compound | Ruptures bacterial cell wall, leading to leakage of contents and cell death May disrupt bacterial metabolic pathways, inhibiting cell growth |
| Chlorhexidine gluconate | Cationic bis-biguanide | Ruptures bacterial cell wall, leading to leakage of contents and cell death Binds to salivary mucins and oral surfaces, which inhibits bacterial colonization Binds to bacteria, inhibiting their adsorption onto teeth surfaces Penetrates the plaque biofilm to exert antimicrobial effects |

Table 1. Descriptions and mechanisms of action of the three most-commonly used active ingredients in antimicrobial mouthrinses.^{21,22}

By virtue of their various mechanisms of action, chlorhexidine gluconate, essential oils, and cetylpyridinium chloride exhibit broad spectrums of antimicrobial activity, including activity against Gram positive and Gram negative bacteria, and against a variety of aerobic and anaerobic bacteria.²¹ The antimicrobial activity of essential oils is complemented by their anti-inflammatory activity, which is achieved via inhibition of prostaglandin synthetase, an enzyme involved in the formation of prostaglandin inflammatory mediators.²¹

An additional benefit of an essential oils-containing mouthrinse is that it has a neutral electrical charge and therefore does not interact with charged ions found in dentifrices, such as sodium lauryl sulphate.^{21,22} Both chlorhexidine gluconate and cetylpyridinium chloride are cationic rinses and susceptible to this interaction, which may lead to reduced biological activity. Also, unlike chlorhexidine gluconate, the essential oils are not inhibited by blood proteins, suggesting an effective user-friendly option in conjunction with mechanical brushing and interproximal cleaning.²¹

Clinical efficacy

Two systematic reviews of published evidence support the effectiveness of antimicrobial mouthrinses in reducing plaque and gingivitis when used as an adjunct to home care.^{22,23} The majority of studies have shown that daily use of mouthrinses containing chlorhexidine gluconate or essential oils provide clinically significant anti-gingivitis and anti-plaque benefits compared with inactive control mouthrinse. Mouthrinse containing cetylpyridinium chloride appeared to provide more limited clinical benefits, possibly due to fewer clinical trials evaluating the same formulations of cetylpyridinium chloride. In addition to their anti-plaque and anti-gingivitis effects, the majority of antimicrobial mouthrinses have shown beneficial effects in reducing oral malodour in both short- and longer-term studies.^{24,25}

In a meta-analysis of head-to-head studies that evaluated the effects of long-term (≥ 4 weeks) use of chlorhexidine gluconate- versus essential oils-based mouthrinses, chlorhexidine gluconate produced better results for anti-plaque and anti-gingivitis benefits but was associated with considerably more staining and calculus.²⁶ For the long-term control of gingival inflammation, both active ingredients produced similar results. Based on these findings, the investigators concluded that an essential oils-containing mouthrinse is a reliable alternative to a chlorhexidine gluconate-containing mouthrinse where long-term anti-inflammatory oral care is deemed beneficial. Indeed, a recent meta-analysis of long-term (6-month) clinical trials supports the benefits of daily essential oils-based mouthrinse use for gingivitis and plaque control beyond that of mechanical oral hygiene alone.²⁷ For indications where plaque control is the main focus, chlorhexidine gluconate remains the active ingredient of first choice.²⁶

PRACTICE TIPS 2 – PATIENT ORAL HYGIENE

1. Remind patients that mouthrinses are not a replacement for mechanical oral hygiene practices – rinses are an adjunct to professional and home mechanical hygiene practices.
2. Brushing and flossing are the primary means of removing plaque at home – adjunctive use of an antimicrobial mouthrinse helps to reduce plaque build-up and gingivitis.
3. Advise patients to choose a mouthrinse with a pleasant usage experience – enjoyment using the product will increase adherence.
4. Advise patients to follow usage instructions; in particular, to not dilute mouthrinses or rinse for less than the recommended amount of time since doing so may reduce their effectiveness.
5. Advise patients not to smoke and, where possible, facilitate access to a smoking cessation programme.
6. Smoking cessation guidelines and training programmes for oral health professionals are available and should be used.

Safety and tolerability

Studies show that daily, long-term use of chlorhexidine gluconate or essential oils mouthrinses does not adversely affect oral microbial flora, including no microbial overgrowth, opportunistic infection, or development of microbial resistance.^{21,22} Long-term use of chlorhexidine gluconate-, essential oils-, or cetylpyridinium chloride-containing mouthrinses does not appear to contribute to the development of soft tissue lesions or mucosal aberrations.²¹ However, taste perception alteration, increased supragingival calculus formation and brown staining of the teeth and other oral surfaces is associated with the use of mouthrinses containing chlorhexidine gluconate.^{21,26} In some cases the staining is severe, requiring professional prophylaxis.²⁸

Regarding concerns that use of alcohol-based mouthrinses can result in desiccation of the oral mucosa, leading to xerostomia, clinical studies have shown no significant difference in salivary flow rate with alcohol-based mouthrinse.^{21,22,29} Importantly, current evidence does not indicate a causal link between the use of alcohol-based mouthrinses and the risk of oral and pharyngeal cancer.^{21,29,30}

Smoking and oral health

Tobacco smoking is a major factor associated with chronic periodontal disease and contributes to higher levels of tooth, attachment, and bone loss.³¹⁻³⁵ For example, a large cross-sectional study of Thai men who were smokers, identified tobacco smoking as being directly associated with periodontitis and indirectly associated with tooth loss.³⁴

Pathology

The mechanisms behind the destructive effects of smoking on the periodontal tissues are not fully understood but are likely to involve interference with vascular and inflammatory processes and the negative effects of nicotine and carbon monoxide in tobacco smoke on mucosal healing.^{36,37} Smoking modifies neutrophil function, which can lead to a shift to a more pathogenic oral microbiome, causes sustained peripheral vasoconstriction leading to a decrease in gingival blood flow, moderates immunoinflammatory responses, delays the healing potential of periodontal tissues, and increases the potential for bone loss.³⁷ Additionally, there appears to be a cumulative effect of smoking on attachment loss.³⁵

Smoking cessation

The risk of periodontal disease has been estimated to be 3- to 20-fold higher in smokers than in non- or never-smokers.^{31,32,38} The rate of progression of periodontal disease is increased in smokers, but reverts to that of a non-smoker following smoking cessation.³⁸ Indeed, there is evidence indicating that smoking cessation is an important component of periodontal treatment, and smokers should be encouraged to quit as part of their overall oral health maintenance.³⁹

The incidence of oral cancer, specifically squamous cell carcinoma, is four to seven times greater in smokers compared with non-smokers,⁴⁰ and when considering the associated increased periodontal disease morbidity and poor wound healing, smoking cessation counselling and support should form an essential role of all dental practitioners.⁴¹

A Cochrane review determined that, based on available clinical trial evidence, behavioural interventions for tobacco cessation conducted by oral health professionals, incorporating an oral examination component in the dental office or community setting, may increase rates of smoking cessation.⁴⁰ UK researchers, who demonstrated that quit rates following smoking cessation advice given as part of a periodontal treatment compare favourably with national quit rates achieved in specialist smoking cessation clinics, concluded that the dental profession has a crucial role to play in smoking cessation for patients with chronic periodontitis.⁴²

Against this background, the American Dental Association and the Australian Dental Association advise that dental healthcare professionals should be encouraged to educate the public on the adverse health implications of smoking as well as how to quit, and that appropriate smoking cessation programmes should be integrated into dental practices.^{43,44} Similarly, Singapore's Health Promotion Board (with endorsement from the Singapore Dental Association) states in its clinical practice guidelines on treating tobacco use and dependence that every healthcare professional should make treating tobacco dependence a priority during a patient consultation.⁴⁵ Online smoking cessation guidelines and training programmes for dental professionals are available (links to a selection of these are provided in the associated **Highlights Box**).

Of note, motivational interviewing techniques are encouraged along with nicotine replacement therapy to double the chances of long-term quitting.^{41,46} In Indonesia, support for smoking cessation therapies (as one of multiple tobacco harm-reduction strategies) has been advocated in the absence of national policy guidelines on tobacco harm reduction.⁴⁷

Smoking Cessation Guidelines and Training Programmes for Dental Professionals

TRAINING PROGRAMMES

Tobacco Control Office, Department of Health, Hong Kong SAR Government
eLearning ABC - Help your client quit smoking
https://www.tco.gov.hk/english/quitting/elearning_abc.html

Singapore Health Promotions Board - Ministry of Health
Smoking Cessation Consultant Training
<https://www.hpb.gov.sg/community/smoking-cessation-programme>

GUIDELINES

FDI World Dental Federation (FDI) and the World Health Organization (WHO):
Tobacco or oral health: An advocacy guide for oral health professionals
http://www.who.int/oral_health/media/orh_tobacco_fdi_book.pdf

Philippines Department of Health:
Smoking Cessation Programme
<http://portal.doh.gov.ph/content/smoking-cessation-program-0.html>

Royal Australian College of General Practitioners:
Supporting Smoking Cessation: A Guide for Health Professionals
<http://www.racgp.org.au/your-practice/guidelines/smoking-cessation/>

Tobacco Control Office, Department of Health, Hong Kong SAR Government
Smoking Cessation Information Kit
<https://www.tco.gov.hk/english/quitting/files/kit15.pdf>

US National Institutes of Health:
Tobacco Control Monograph Series
<https://cancercontrol.cancer.gov/brp/tcrb/monographs/>

RESEARCH REVIEW EDUCATIONAL SERIES

Antimicrobial Mouthrinse in Dental Practice and Home Oral Hygiene Smoking Cessation and Oral Health



EXPERT COMMENTARY BY VIVIAN VIRATA (PHILIPPINES)

Antimicrobial mouthrinse is an essential tool for infection control in a dental office. As a pre-procedural rinse, it reduces bacterial aerosols prior to any dental procedure. Also, as an adjunct to oral home care, mouth rinsing will be able to "hard to reach" areas in the oral cavity, thereby controlling bacteria and delaying the onset of the disease process.

The four essential oils-containing mouthrinse has extensive studies demonstrating its efficacy, including anti-plaque and gingival index's scores, and antibacterial activity.

Smoking cessation for 10 years will greatly help treatment outcomes, almost the same as never smokers. However, the use of a mouthrinse will not negate the destructive effects of smoking.

EXPERT COMMENTARY BY EDWARD LO (HONG KONG)

This article nicely summarizes the important information on infection control in dental practice and on antimicrobial mouthrinse. It also draws attention to the essential role of the dentist in promoting smoking cessation among their patients.

Implementation of proper infection control measures in the dental clinic is of vital importance in the Southeast Asian region where communicable diseases are common and their spread needs tight control in the populous areas. A fundamental principle in providing healthcare is to minimize avoidable harm or health risk to the service users. Dentists should take up the responsibility to maintain a clean environment and to ensure minimal risk of cross-infection in their clinics.

Antimicrobial mouthrinse, as an adjunct to normal mechanical oral hygiene measures, has been shown to be effective in reducing dental plaque and gingivitis among different population groups. However, its role in promoting oral health is not commonly highlighted

in oral health education activities and is probably under-utilized by the people in most countries in the region. Dental professionals need to have a better understanding of the different antimicrobial mouthrinses and help the public to make an informed choice.

It is well-known that use of tobacco, such as smoking, is associated with many serious diseases and health problems, e.g. lung cancer and cardiovascular diseases. Its negative impact on oral health has also been shown. Using a common risk factor approach in health promotion is advocated by the World Health Organization and the health authorities in many countries in the region. Dentists, as key members of the health profession and leaders of the oral healthcare team, should play an active role in promoting smoking cessation. This article contains links to many good sources of training and guidelines in this aspect and is very helpful for the dentists in the region.

TAKE-HOME MESSAGES:

- Aerosols and splatter are a potential source of cross-infection in the dental surgery.
- Steps should be taken to minimize the generation of aerosols and splatter, e.g. use of high-evacuation, rubber dams, and pre-procedural use of antimicrobial mouthrinse.
- As an adjunct to mechanical plaque removal, the daily use of an antimicrobial mouthrinse helps to reduce plaque formation and gingivitis.
- Smoking is a risk factor for periodontal disease and dental health practitioners should encourage smoking cessation and facilitate access to a smoking cessation programme, where available.

REFERENCES

- Centers for Disease Control and Prevention. Guidelines for infection control in dental health-care settings - 2003. Atlanta, GA: Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion. 2003. Available from: <https://www.cdc.gov/mmwr/preview/mmwrhtml/rr5217a1.htm>.
- Centers for Disease Control and Prevention. Summary of infection prevention practices in dental settings: Basic expectations for safe care. Atlanta, GA: Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion. 2016. Available from: <https://www.cdc.gov/oralhealth/infectioncontrol/pdf/safe-care.pdf>.
- Bird DL. Chapter 19. Disease transmission and infection prevention. In: Bird DL, et al., editors. Student Workbook for Modern Dental Assisting. Vol. 11th ed. New York, NY: Elsevier Saunders. 2014: 83-100.
- Day CJ, et al. Aerosols and splatter in dentistry--a neglected menace? Dent Update. 2006;33(10):601-2, 4-6.
- Rautema R, et al. Bacterial aerosols in dental practice - a potential hospital infection problem? J Hosp Infect. 2006;64(1):76-81.
- Harrel SK, et al. Aerosols and splatter in dentistry: a brief review of the literature and infection control implications. J Am Dent Assoc. 2004;135(4):429-37.
- Australian Dental Association (ADA). ADA's guidelines for infection control. 2015. St Leonards, NSW: Australian Dental Association. Available from: <http://adawa.com.au/media/6-ADA-Guidelines-for-Infection-Control-Third-Edition.pdf>.
- Puttaiah R, et al. Comparison of knowledge, attitudes and practice of dental safety from eight countries at the turn of the century. J Contemp Dent Pract. 2011;12(1):1-7.
- Oosthuysen J, et al. Compliance with infection prevention and control in oral health-care facilities: a global perspective. Int Dent J. 2014;64(6):297-311.
- DePaola LG, et al. Effect of an antiseptic mouthrinse on salivary microbiota. Am J Dent. 1996;9(3):93-5.
- Dewker NR, et al. A study to evaluate and compare the efficacy of preprocedural mouthrinsing and high volume evacuator attachment alone and in combination in reducing the amount of viable aerosols produced during ultrasonic scaling procedure. J Contemp Dent Pract. 2012;13(5):681-9.
- Feres M, et al. The effectiveness of a preprocedural mouthrinse containing cetylpyridinium chloride in reducing bacteria in the dental office. J Am Dent Assoc. 2010;141(4):415-22.
- Fine DH, et al. Reduction of viable bacteria in dental aerosols by preprocedural rinsing with an antiseptic mouthrinse. Am J Dent. 1993;6(5):219-21.
- Fine DH, et al. Efficacy of preprocedural rinsing with an antiseptic in reducing viable bacteria in dental aerosols. J Periodontol. 1992;63(10):821-4.
- Gupta G, et al. Efficacy of preprocedural mouth rinsing in reducing aerosol contamination produced by ultrasonic scaler: a pilot study. J Periodontol. 2014;85(4):562-8.
- Shetty SK, et al. Compare the efficacy of two commercially available mouthrinses in reducing viable bacterial count in dental aerosol produced during ultrasonic scaling when used as a preprocedural rinse. J Contemp Dent Pract. 2013;14(5):848-51.
- Lovegrove JM. Dental plaque revisited: bacteria associated with periodontal disease. J N Z Soc Periodontol. 2004;8(7):7-21.
- Zijge V, et al. Oral biofilm architecture on natural teeth. PLoS One. 2010;5(2):e9321.
- Arweiler NB, et al. The Oral Microbiota. Adv Exp Med Biol. 2016;902:45-60.
- Canro LP, et al. The expected effect on oral health of dental plaque control through mechanical removal. Periodontol 2000. 1993;8:60-74.
- DePaola LG, et al. Safety and efficacy of antimicrobial mouthrinses in clinical practice. J Dent Hyg. 2007(Special supplement):13-25.
- Ossó D, et al. Antiseptic mouth rinses: an update on comparative effectiveness, risks and recommendations. J Dent Hyg. 2013;87(1):10-8.
- Gunsolley JC. Clinical efficacy of antimicrobial mouthrinses. J Dent. 2010;38 Suppl 1:S6-10.
- Blom T, et al. The effect of mouthrinses on oral malodor: a systematic review. Int J Dent Hyg. 2012;10(3):209-22.
- Fedorowicz Z, et al. Mouthrinses for the treatment of halitosis. Cochrane Database Syst Rev. 2008(4):Cd006701.
- Van Leeuwen MP, et al. Essential oils compared to chlorhexidine with respect to plaque and parameters of gingival inflammation: a systematic review. J Periodontol. 2011;82(2):174-94.
- Araujo MW, et al. Meta-analysis of the effect of an essential oil-containing mouthrinse on gingivitis and plaque. J Am Dent Assoc. 2015;146(8):610-22.
- Adams D, et al. Mouthrinses. Adv Dent Res. 1994;8(2):291-301.
- Fischman SL, et al. Use of essential oil-containing mouthrinses by xerostomic individuals: determination of potential for oral mucosal irritation. Am J Dent. 2004;17(1):23-6.
- La Vecchia C. Mouthwash and oral cancer risk: an update. Oral Oncol. 2009;45(3):198-200.
- Bergstrom J. Tobacco smoking and chronic destructive periodontal disease. Odontology. 2004;92(1):1-8.
- Cesar Neto JB, et al. Smoking and periodontal tissues: a review. Braz Oral Res. 2012;26 Suppl 1:25-31.
- Razali M, et al. A retrospective study of periodontal disease severity in smokers and non-smokers. Br Dent J. 2005;198(8):495-8; discussion 85.
- Chatrchaiwivatana S, et al. Periodontitis associated with tobacco smoking among rural Khon Kaen Thai males: analysis of two data sets. J Med Assoc Thai. 2009;92(11):1524-31.
- Zeng J, et al. Reexamining the association between smoking and periodontitis in the dunedin study with an enhanced analytical approach. J Periodontol. 2014;85(10):1390-7.
- Lee J, et al. Cigarette smoking and inflammation: cellular and molecular mechanisms. J Dent Res. 2012;91(2):142-9.
- Ojima M, et al. Destructive effects of smoking on molecular and genetic factors of periodontal disease. Tob Induc Dis. 2010;8:4.
- Johnson GK, et al. Cigarette smoking and the periodontal patient. J Periodontol. 2004;75(2):196-209.
- Chambone L, et al. Effects of smoking cessation on the outcomes of non-surgical periodontal therapy: a systematic review and individual patient data meta-analysis. J Clin Periodontol. 2013;40(6):607-15.
- Carr AB, et al. Interventions for tobacco cessation in the dental setting. Cochrane Database Syst Rev. 2012(6):Cd005084.
- FDI World Dental Federation (FDI) and the World Health Organization (WHO). Tobacco or oral health: An advocacy guide for oral health professionals. Lowestoft, UK: FDI World Dental Press Ltd. 2005.
- Nasry HA, et al. Smoking cessation advice for patients with chronic periodontitis. Br Dent J. 2006;200(5):272-5; discussion 65.
- Policies and recommendations on tobacco use: Dentist's role in preventing tobacco use. Chicago, IL: ADA American Dental Association. Last update date: 23/01/17. Available from: <http://www.ada.org/en/about-the-ada/ada-positions-policies-and-statements/policies-and-recommendations-on-tobacco-use>. [Date accessed: 23/02/17].
- Australian Dental Association Policies. Policy statement 2.2.4 - community oral health promotion: Tobacco. St Leonards: Australian Dental Association Inc., 2014. Available from: http://www.ada.org.au/Dental-Professionals/Policies/National-Oral-Health-2-2-4-Tobacco/ADAPolicies_2-2-4_Tobacco_V1.
- Chan K, et al. Health promotion board-ministry of health clinical practice guidelines: treating tobacco use and dependence. Singapore Med J. 2013;54(7):411-5; quiz 6.
- Royal Australian College of General Practitioners. Supporting smoking cessation: A guide for health professionals. Melbourne, VA: Royal Australian College of General Practitioners (RACGP). 2014. Available from: <http://www.racgp.org.au/your-practice/guidelines/smoking-cessation/>.
- Nurwidya F, et al. Strategies for an effective tobacco harm reduction policy in Indonesia. Epidemiol Health. 2014;36:e2014035.

SUBSCRIBE FREE OF CHARGE AT RESEARCH REVIEW www.researchreview.com/Philippines

Johnson & Johnson

This publication has been created with an educational grant from Johnson and Johnson Asia Pacific. The content is entirely independent and based on published studies and the writer and commentators' opinions.